

**Appraisal of polycyclic aromatic hydrocarbons (PAHs) in environmental matrices
by analytical pyrolysis (Py-GC/MS)**

J.A. González-Pérez^{1*}, G. Almendros², J.M. de la Rosa¹, F.J. González-Vila¹

1) IRNAS-CSIC, Av. Reina Mercedes, 10, 41012-Seville, Spain

2) MNCN-CSIC, c/Serrano 115-B, 41080-Madrid, Spain

*Corresponding author:

Tel +34954624711; Fax +34 954644002; e-mail jag@irnase.csic.es

Abstract

Large number of studies has demonstrated the usefulness of flash pyrolysis in association with GC-MS in releasing diagnostic volatile compounds informing about the complex composition of natural organic matter (NOM). This knowledge is of interest as regards monitoring agricultural and environmental processes including pollution. In this context the term NOM comprises a wide array of biomacromolecules (lignins, polysaccharides, proteins, lipid polymers, etc.), as well as complex three-dimensional macromolecules, either labile or recalcitrant, and which are referred to with generic terms such as Dissolved Organic Matter, Humic Substances, Black Carbon, etc. In this communication an updated overview on recent advances achieved by Analytical Pyrolysis in the structural characterization of recalcitrant/fossil/ NOM is presented, with particular focus in the detection of polycyclic aromatic compounds (PAHs) within complex organic matrices.

Keywords: Black carbon, Burning, Fire, Humic substances, Lignite, Pyrogenic organic matter, Soil organic matter

Introduction

Pyrolysis is the breaking apart of large molecules into smaller ones using thermal energy in the absence of oxygen. The combination of pyrolysis with separation and detection means is known as analytical pyrolysis. This technique is a powerful tool widely used for the direct characterization of organic mixtures with diverse origin which, owing to their complexity, are difficult to analyse by both conventional destructive or non-destructive methods. Pyrolysis is routinely used as a forensic tool to fingerprint or trace specific analytes within complex substances such as tyre rubber, textiles, paint, glue, paper coatings, petrochemical sources, or plant-derived materials including wood, coal and fossil remains, archaeological and cultural heritage artefacts, bacteria, and the whole range of synthetic polymers [1–5].

Pyrolysis of NOM generates a wide range of products with diverse structure that can be related to their biochemical origin. In particular, specific pyrolytic compounds from vascular plant debris are often monitored as tracers of the fate of the source material [6]. In the case of soil or sedimentary organic matter, analytical pyrolysis has enlightened crucial structural aspects in a wide range of substances from kerogens or peats to humic-like substances from composted organic matter [7–10], including soil organic matter [11–14], sedimentary organic matter [15–17], pyrogenic C-forms i.e., black carbon (BC) and other highly refractory forms of organic matter accumulated and transformed in soils and sediments, or newly-formed during forest fires [18–28].

This is the case with BC, which is understood as a continuum from partly charred plant material, through char and charcoal to graphite and soot particles condensed from the

gas phase [29–31]. As biomass is charred, aryl and *O*-aryl structures (aromaticity) increase in concentration with a concomitant loss in carbohydrate and lignin structures [32]. Micromorphological and chemical features of this refractory material are described elsewhere [33–36]. Black carbon behaves as a highly aromatic material and, in the case of BC from vegetation fires [37], is considered to consist of randomly oriented stacks of few graphitic layers. In other cases BC is not necessarily graphitic, and this black material, even when completely amorphous, may contain both a substantial alkyl domain and oxygen content (endocyclic or lactone-like) structures [34, 36, 38]. Black carbon takes part in a variety of relevant environmental processes. Thus, it affects Earth climate through its light-absorbing effect, but it is also thought that BC may be part of the missing carbon in the global carbon balance [39–40 and references therein] as well as a carrier-phase of pollutants playing a key role in the sorption of contaminants including PAHs [41]. Although the molecular composition and source-dependent structural variability of BC remain largely unknown, it is admitted that PAHs are its most typical discrete components.

Polycyclic aromatic hydrocarbons (PAHs) are ubiquitous components of natural organic matter subjected to fire's impact or geothermal conditions. There are three major environmental sources of PAHs: combustion of organic matter (pyrogenic), petroleum (petrogenic) and natural sources including volcanism [42 and references therein]. While local areas can be impacted by petrogenic PAHs, combustion of organic matter is the largest source of PAHs to the environment with biogenic sources of PAHs limited to only a few compounds (i.e. retene and perylene). In general it is considered that pyrolytic PAHs include all parent compounds other than retene and perylene, having MW > 178. Alkylated homologue PAHs usually derive from fossil sources [43–45]. In

any case, and as regards PAHs found in environmental matrices like soils, sediments or BC we could consider their origin either as native constituents or incorporated by sorption of allochthonous pyrogenic or anthropogenic products.

The environmental occurrence of PAHs is frequently underestimated by both wet chemical degradation or spectroscopic approaches, which often lead to patterns which are often meaningless from the structural viewpoint in which heavily polycondensed aromatic material often represent an ‘invisible’ domain, either by its frequent spectroscopic underestimation or the nonstoichiometric yields after chemical degradation [46]. In fact, frequent routine methodological approaches to study PAHs from geological samples are based on analyses of solvent extracts and further chromatographic separation [47–49]. However, detection of PAHs by direct pyrolysis of geological materials is feasible and has been reported [50–53], evidencing a series of advantages for the rapid examination of PAHs in whole sediments.

The main aim of this review study is to examine antecedents and the reliability of analytical pyrolysis techniques in evidencing the occurrence of PAH compounds in complex organic matrices. In addition, the potential of Py-GC/MS as analytical technique to detect PAHs in environmental samples was tested in a set of standard samples that has been thoroughly characterized by previous studies using non-destructive (e.g., ^{13}C NMR spectroscopy) techniques. These included charred biomass, soils, sediments, refractory organic matter (BC), soot, coal and lignite. PAH-lacking samples were used as negative controls.

Capability of pyrolysis tools and conditions for PAH analysis

Analytical pyrolysis has been used as a tool for the analysis of PAHs included in environmental matrices (i.e soil and sediments), as pollutants, constituents (i.e. coals and black carbon) or pyrogenic (i.e. chars and charred materials from forest fires). Medina-Vera [50] used a platinum coil pyrolyser to screen PAHs in pure standard mixtures and sediment samples. Adjusting the analytical conditions the author was able to thermally desorb (350 °C) the PAHs from sediment samples without any pre-treatment. The method was proved to be appropriate for the detection of compounds ranging from naphthalene to pyrene, which demonstrated that analytical pyrolysis is a suitable and convenient technique for PAHs screening in sediment samples [50]. A modification of this technique was developed by Faure and coworkers as a rapid contamination screening tool. The so called thermodesorption GC/MS (TdGC/MS) was found efficient for the rapid assessment of sediment contamination by free pollutants (especially by PAHs) with no interferences from possible cyclization and aromatization artifacts formed in the presence of clays in the media. It was also considered this technique as a valuable tool to provide crucial information to environmental agencies for long term management of rivers and sediments [51–52].

Terán et al. [16] used double shot analytical pyrolysis in a screening of organic contaminants in sediments from the South Iberian Peninsula detecting series of linear alkylbenzenes (most probably from detergents), PAHs (petroleum or combustion products) and alkylnitriles with a probable origin in the industry and urban anthropogenic activities. They also established optimum pyrolysis temperature for pollution detection of these contaminants and demonstrated the usefulness of Py–GC/MS as a tool for rapid screening of organic contaminants in sediments. The same pyrolysis technique was also a very useful technique for the on-line determination of

PAHs in airborne particulate matter (soot/BC) trapped on cellulose filters without the need of using any sample pretreatment [53].

It has been found that Py-GC/MS pattern of coals is highly consistent with the chemical structure as inferred from NMR and elemental composition data. In fact the abundance of aromatics in the pyrolysates increased with coal rank in such a way that this could be used as coal rank parameter [54].

Curie point analytical pyrolysis and thermally assisted hydrolysis and methylation (THM) has been used to detect aged black carbon (BC) in soils. Kaal et al. [27, 28, 31, 37, 40] first studied ancient fire-affected organic matter thorough analytical pyrolysis and found a large proportion of PAHs, benzenes and benzonitriles in the pyrolysates of extractable soil organic matter derived from BC. They concluded that BC in soil conforms a partially oxidised, highly aromatic but weakly condensed network, containing significant amounts of N as previously suggested [55].

Recently González-Pérez et al. [56] used double shot pyrolysis to investigate organic matter evolution in lignite-rich acid mine soils. Apart from the normal markers produced by the recent OM, a wide range of PAHs were detected. These include natural PAHs like retene, considered a diagenetic product of dehydroabietic acid and a typical geochemical marker for lignite [57], and series of up to four benzene ring structures (pyrene) in the sediments. Traces of PAHs of up to five benzene rings were found in the Miocene lignite pyrolysates. Also major series of two-ring compounds mainly alkylnaphthalenes in addition to dibenzofurans and fluorene derivatives were found. The authors concluded that compounds present in the lignite substrate but lacking in the

vegetation pyrolysates i.e. alkylbenzene series, 1H-inden-1-one, 2,3-dihydro-3,3,4,5-tetramethyl-, as well as PAHs other than retene, were appropriate markers betraying the contribution from lignite to the soil/sediment organic matter.

Py-GC/MS of standard samples

Reference samples studied

The samples studied are indicated in Table 1 and consists of:

A) BC-rich materials; chars (chestnut wood and rice grass), obtained from dried biomass in an electric oven at 400 °C for 5 h under inert atmosphere [58], a soot and a low bituminous coal consisting of *c.* 10% pyrogenic inertinite, the characteristics of which being described elsewhere [59].

B) Environmental matrices including polyaromatic hydrocarbons. A Cretaceous sediment (Barremian-Albian; 130–100 Million years, dated by micropalaeontological data) was studied, this corresponds to the Cerrajón Formation (South of Jaén, Spain) in the Baetic Cordillera. This stratigraphic unit is a shale composed mainly of silty marls with organic matter interbedded with several layers of sandstones [60–61]. In addition, three soils thought to contain PAHs were included in this group; two black isohumic soils (Vertisol and a Chernozem) [59] and a Rendolic soil from ‘Sierra de Cazorla’ (Southern Spain) affected or not by a forest fire [12] that is a frequent source of BC in Mediterranean environments.

C) Two black-coloured organic samples used as negative controls: a melanoidin produced from urea + glucose at 90 °C for 30 days and a marine sediment (oxidation-

resistant organic matter fraction: ROM) from offshore Huelva (SW Spain) coast. As in the above case, and despite its dark colour, after analytical characterization by ^{13}C NMR no appreciable aromatic carbons were detected with the sample consisting of predominantly non-hydrolysable material with strong aliphatic nature [59, 62].

Conditions of the Py-GC/MS experiments

Pyrolysis-gas chromatography–mass spectrometry (Py-GC/MS) was performed using a double-shot pyrolyzer (Frontier Laboratories, model 2020i) attached to a GC/MS system Agilent 6890N. Unaltered ground samples (5–50 mg depending on C content) were placed in small crucible capsules and introduced into the furnace, which was preheated at 500 °C for 1 min. In certain experiments the final temperature was 800 °C and in this case it is detailed in the corresponding Figure caption. The evolved gases were then directly injected into the gas chromatograph-mass spectrometer for analysis. The gas chromatograph was equipped with a low-to-mid polarity-fused silica capillary column (J&W Scientific) of 30 m \times 250 μm \times 0.25 μm film thickness (Ref. DB-1701). The carrier gas was helium at a controlled flow of 1 mL min $^{-1}$. The detector consisted of an Agilent 5973 mass selective detector, and mass spectra were acquired at 70 eV ionizing energy. Compound assignment was achieved via single-ion monitoring for various homologous series, via low-resolution mass spectrometry, and via comparison with published and stored (NIST and Wiley libraries) data. Traces corresponding to selected homologous series of chemical families were obtained by single ion monitoring (SIM) of characteristic ions.

In order to validate the reliability of the Py-GC/MS approach to identify PAHs in a large variety of reference materials, the results of a series of selected independent experiments are described and discussed below.

Release of PAHs from BC reference materials

The pyrograms from BC-containing materials (lignocellulose-derived chars from chestnut wood and rice grass), Vertisol and Chernozem soils, bituminous coal, urban soot and the negative control reference material (melanoidin) are shown in Fig. 1. These samples correspond to reference materials studied in a previous comprehensive interlaboratory experiment aiming to quantify fire-derived carbon in soils and sediments [59]. The results evidence that, with limitations inherent to the chromatographic separations, analytical pyrolysis was successful in identifying a wide range of aromatic hydrocarbons presumably released from the BC moiety including PAHs up to six benzene rings (e.g. benzoperylene, C₂₂). No PAH, even at trace amount, was detected in the negative control.

The pyrograms of oxidation-resistant organic matter from the coast of Huelva (SW Spain) [62] reveals that this material has a strongly aliphatic nature, yielding *n*-alkanes and *n*-fatty acids as major pyrolysis compounds whereas aromatics were practically lacking, as corresponds to a marine origin. On the other side, the major pyrolysis compounds from lignocellulose char consisted mainly of aromatic and heterocyclic compounds like benzofurans that are well known pyrolysis products from incompletely charred lignocellulosic materials/coal tars [63] in addition to a major levoglucosan peak arising from the pyrolysis of carbohydrates in celluloses [64]. Solid-state carbon nuclear

magnetic resonance (^{13}C -CP/MAS-NMR) of the same samples complements [27] and, in this case, corroborates the pyrolysis results: this marine refractory material displays a main alkyl-C signal (45–0 ppm) whereas the spectrum from the lignocellulose char displayed a major peak in the aromatic C range (160–110 ppm).

Release of PAHs from Cretaceous sediments

The occurrence of PAHs in both recent and ancient sediments has been extensively reported. In fact, some individual compounds as well as PAH ratios are currently used as biogeochemical proxies for sediment characterization and palaeoclimate assessment [49, 65]. In this third experiment, a Cretaceous sediments is studied by direct Py-GC/MS.

The sediments released a wide set of PAHs including unsubstituted and alkylated homologues of phenanthrene, fluorene, anthracene, pyrene, perylene and coronene (Fig 3). The dominance of cata-condensed structures (i.e. benzo[*e*]pyrene, benzo[*ghi*]fluoranthene) over peri-condensed (i.e. benzo[*a*]pyrene, cyclopenta[*cd*]pyrene) is evident, suggesting a pyrolytic origin at high temperatures. Also, the observation that non-alkylated structures derived from phenantrene, dibenzothiophene and fluoranthene are predominant over their alkylated homologues (Fig 3) indicates a contribution of pyrogenic material to the organic matter in the sediment.

The PAHs distribution found from the Andalusian shale is similar to that previously reported for sites located far from pollution sources (i.e. high mountain lakes and deep

sea sediments), clearly pointing to an origin from natural pyrogenic material. The PAHs pattern found was most probably the result of in situ combustions and/or deposition of soot from forest fires during the frequent fire events in the Cretaceous [66].

Release of PAHs from wildfire-affected soils

Mediterranean forest is highly vulnerable to fire during the dry, hot summer months where temperatures often exceed 40 °C [67]. Forest wildfires contribute to the periodic destruction of Mediterranean and Atlantic ecosystems in the Iberian Peninsula and are considered the main disturbance in the Mediterranean basin [68–70].

It has been indicated that fires' impact in Mediterranean and Atlantic ecosystems may be recognized by severe transformations of soil physical, chemical and biological characteristics associated to qualitative and quantitative changes in the most functional fraction of soil such as organic matter including biota [19, 71–72]. These changes may negatively affect soil health and quality, favouring the occurrence of erosive processes [73]. The assessment of the effects of forest fires on organic matter is one of the most complex subjects in the systematic research of environmental impacts [74]. The identification of biogeochemical markers for the occurrence of forest fire events, as well as to monitor ecosystems recovery is important for planning environmental restoration actions after wildfires [12]. Among the obvious candidates as tracers for the occurrence of wildfires in soils are pyrogenic PAHs.

For this case study the identification of PAHs in soil was assessed in a comparative manner between the fire-affected soil and the neighbour, non-affected one, with the same pedological and physiographical characteristics. When comparing the ion traces

for diagnostic PAHs in whole soil pyrograms, structures from one to four or more aromatic rings were observed only in the fire-affected soil, being traceable down in the soil profile (0 to 20 cm depth) (Fig. 4). This fact, in addition to the higher relative abundance of three and four benzene rings PAHs observed in soil top layers of the fire-affected soil, indicates a strong modification of the soil organic matter, either directly by fire or by charred material inputs additions from the burned vegetation. The effect was clearer in the uppermost soil layers.

Conclusions

Usual methodologies to analyse PAHs are in general time-consuming, and have additional drawbacks such as the need of previous extraction steps with organic solvents. Pyrolysis in association with GC/MS has shown valuable potential in the structural characterization of many different biopolymers and geomacromolecules, evidencing PAHs series of up to seven benzene rings from environmental chemically complex matrices. These conspicuous series of PAHs were not detected in other recalcitrant natural substrates where ^{13}C NMR indicates an almost completely aliphatic nature. Although not a quantitative technique, analytical pyrolysis (Py-GC/MS) is proposed as a fast alternative approach to detect PAHs in environmental samples. Amongst the advantages are: minimum sample handling steps, thus precluding selective losses or sample contamination, speed of analysis and the need of small amounts of samples.

300

301 **Acknowledgements**

302 This work has been partly funded by the European Union RECARE (FP7-ENV-2013
303 Grant Agreement No 603498) and the Spanish ‘Ministerio de Economía y
304 Competitividad’ through project GEOFIRE (CGL2012-38655-C04-01). Prof. Heike K
305 Knicker is acknowledged for the NMR analysis. Thanks are also due to Dr. Marianne
306 Blazsó and two anonymous reviewers for their always helpful comments while writing
307 this MS.

308

References

- [1] G. Almendros, J. Dorado, F.J. González-Vila, F. Martín, J. Sanz, C. Álvarez-Ramis, L. Stuchlik, Molecular characterization of fossil organic matter in *Glyptostrobus europaeus* remains from the Orawa basin (Poland). Comparison of pyrolytic techniques. *Fuel* 78 (1999) 745–752.
- [2] T.P. Wampler, Introduction to pyrolysis-capillary gas chromatography. *J. Chromatogr. A*. 842 (1999) 207–220.
- [3] M.J. Simpson, B. Chefetz, A.P. Deshmukh, P.G. Hatcher, Comparison of polycyclic aromatic hydrocarbon distributions and sedimentary organic matter characteristics in contaminated, coastal sediments from Pensacola Bay, Florida. *Mar. Environ. Res.* 59 (2005) 139–163.
- [4] J.C. Frade, M.I. Ribeiro, J. Graça, J. Rodrigues, Applying pyrolysis-gas chromatography/mass spectrometry to the identification of oriental lacquers: study of two lacquered shields. *Anal. Bioanal. Chem.* 395 (2009) 2167–2174.
- [5] D. Fabbri, A. Adamiano, C. Torri, GC-MS determination of polycyclic aromatic hydrocarbons evolved from pyrolysis of biomass. *Anal. Bioanal. Chem.* 397 (2010) 309–317.
- [6] J.I. Hedges, F.G. Prahl, Early diagenesis: consequences for applications of molecular biomarkers, In: M. Engel and S.A. Macko (Eds), *Organic Geochemistry* Plenum Press. 1993, pp. 237–253.
- [7] F.J. González-Vila, G. Almendros, P. Tinoco, J. Rodríguez, Nitrogen speciation and pyrolytic patterns of N¹⁵-labelled soil and compost fractions. *J. Anal. Appl. Pyrolysis* 58 (2001) 329–339.

- [8] F.J. González-Vila, J.A. González-Pérez, K. Akdi, M.D. Gómis, F. Pérez-Barrera, T. Verdejo, Assessing the efficiency of urban waste biocomposting by analytical pyrolysis (Py-GC/MS). *Bioresource Technol.* 100 (2009) 1304–1309.
- [9] J.A. González-Pérez, F.J. González-Vila, G. Almendros, M.C. Zancada, O. Polvillo, F. Martín, Preferential accumulation of selectively preserved biomacromolecules in the humus fractions from a peat deposit as seen by analytical pyrolysis and spectroscopic techniques. *J. Anal. Appl. Pyrolysis* 68-69 (2003) 287–298.
- [10] A. Piedra-Buena, G. Almendros, F.J. González-Vila, J.A. González-Pérez, H. Knicker, Transformations in carbon and nitrogen-forms in peat subjected to progressive thermal stress as revealed by analytical pyrolysis. *Chem. Ecol.* 26 (2010) 361–370.
- [11] J.A. González-Pérez, C.D. Arbelo, F.J. González-Vila, A. Rodríguez-Rodríguez, G. Almendros, C.M. Armas, O. Polvillo, Molecular features of organic matter in diagnostic horizons from andosols as seen by analytical pyrolysis. *J. Anal. Appl. Pyrolysis* 80 (2007) 369–382.
- [12] J.A. González-Pérez, F.J. González-Vila, R. González-Vázquez, M.E. Arias, J. Rodríguez, H. Knicker, Use of multiple biogeochemical parameters to monitor the recovery of soils after forest fires. *Org. Geochem.* 39 (2008) 940–944.
- [13] C. Pereira de Assis, J.A. González-Pérez, J.M. de la Rosa, I. Jucksch, E. de Sá Mendonça, F.J. González-Vila, Analytical pyrolysis of humic substances from a Latosol (Typic Hapludox) under different land uses in Minas Gerais, Brazil. *J. Anal. Appl. Pyrolysis* 93 (2012) 120–128.
- [14] C. Rumpel, A. Rodríguez-Rodríguez, J.A. González-Pérez, C. Arbelo, A. Chabbi, N. Nunan, F.J. González-Vila, Contrasting composition of free and mineral-

bound organic matter in top- and subsoil horizons of Andosols. *Biol. Fert. Soils* 48 (2012) 401–411.

- [15] O. Polvillo, J.A. González-Pérez, T. Boski, F.J. González-Vila, Structural features of humic acids from a sedimentary sequence in the Guadiana estuary (Portugal-Spain border). *Org. Geochem.* 40 (2009) 20–28.
- [16] A. Terán, F.J. González-Vila, J.A. González-Pérez, Detection of organic contamination in sediments by double-shot pyrolysis–GC/MS. *Environ. Chem. Lett.* 7 (2009) 301–308.
- [17] J.M. de La Rosa, J.A. González-Pérez, F.J. González-Vila, H. Knicker, M.F. Araújo Molecular composition of sedimentary humic acids from South West Iberian Peninsula. A multi-proxy approach. *Org. Geochem.* 42 (2011) 791–802.
- [18] J.A. González-Pérez, F.J. González-Vila, G. Almendros, H. Knicker, O. Polvillo, F. Salas, J.C. Costa, Wildfire and black carbon production in Andalusian Mediterranean forest. In: D.X. Viegas (Ed.) *Forest Fire Research & Wildland Fire Safety* (CD-ROM) Milipress, Rotterdam, The Netherlands, 2002 (ISBN 90 77017 72 0)
- [19] J.A. González-Pérez, F.J. González-Vila, G. Almendros, H. Knicker, The effect of fire on soil organic matter – a review. *Environ. Int.* 30 (2004) 855–870.
- [20] F.J. González-Vila, G. Almendros, Thermal transformation of soil organic matter by natural fires and laboratory-controlled heatings, In: R. Ikan (Ed.). *Natural and laboratory simulated thermal geochemical processes*. Dordrecht: Kluwer Academic Publishing, 2003. p. 153–200.
- [21] F.J. González-Vila, P. Tinoco, G. Almendros, F. Martín, Pyrolysis-GC-MS analysis of the formation and degradation stages of charred residues from lignocellulosic biomass. *J. Agric. Food. Chem.* 49 (2001) 1128–1131.

- 382 [22] F.J. González-Vila, J.A. González-Pérez, O. Polvillo, G. Almendros, H. Knicker,
383 Nature of refractory forms of organic carbon in soils affected by fires. Pyrolytic and
384 spectroscopic approaches, In: D.X. Viegas (Ed.) Forest Fire Research & Wildland
385 Fire Safety (CD-ROM) Millipress, Rotterdam, The Netherlands, 2002 (ISBN 90
386 77017 72 0)
- 387 [23] K. Quénéa, S. Derenne, F.J. González-Vila, J.A. González-Pérez, A. Mariotti, C.
388 Largeau, Double-shot pyrolysis of the non-hydrolysable organic fraction isolated
389 from a sandy forest soil (Landes de Gascogne, South-West France). Comparison
390 with classical Curie point pyrolysis. J. Anal. Appl. Pyrolysis 76 (2006) 271–279.
- 391 [24] H. Knicker, F.J. González-Vila, O. Polvillo, J.A. González-Pérez, G. Almendros,
392 Fire-induced transformation of C- and N- forms in different soil fractions from a
393 Dystric Cambisol under a Mediterranean pine forest (*Pinus pinaster*). Soil. Biol.
394 Biochem. 37 (2005) 701–718.
- 395 [25] C. Rumpel, J.A. González-Pérez, G. Bardoux, C. Largeau, F.J. González-Vila,
396 C. Valentin, Composition and reactivity of morphologically distinct charred
397 materials left after slash-and-burn practices in agricultural tropical soils. Org.
398 Geochem. 38 (2007) 911–922.
- 399 [26] J.M. de la Rosa, J.A. González-Pérez, R. González-Vázquez, H. Knicker, E.
400 López-Capel, D.A.C. Manning, F.J. González-Vila, Use of pyrolysis/GC–MS
401 combined with thermal analysis to monitor C and N changes in soil organic matter
402 from a Mediterranean fire affected forest. Catena 74 (2008) 296–303.
- 403 [27] J. Kaal, S. Brodsowski, J.A. Baldock, K.G.J. Nierop, A. Martínez-Cortizas,
404 Characterisation of aged black carbon using pyrolysis-GC/MS, thermally assisted
405 hydrolysis and methylation (THM), direct and cross-polarisation ¹³C nuclear

magnetic resonance (DP/CP NMR) and the benzenepolycarboxylic acid (BPCA) method. *Org. Geochem.* 39 (2008) 1415–1426.

[28] J. Kaal, A. Martínez-Cortizas, K.G.J. Nierop, P. Buurman, A detailed pyrolysis-GC/MS analysis of a black carbon-rich acidic colluvial soils (Atlantic Ranker) from NW Spain. *Appl. Geochem.* 23 (2008) 2395–2405.

[29] C.A. Masiello, New directions in black carbon organic geochemistry. *Mar. Chem.* 92 (2004) 201–213.

[30] J.I. Hedges, G. Eglinton, P.G. Hatcher, D.L. Kirchman, C. Arnosti, S. Derenne, R.P. Evershed, I. Kögel-Knabner, J.W. de Leeuw, R. Littke, W. Michaelis, J. Rullkötter, The molecularly-uncharacterized component of nonliving organic matter in natural environments. *Org. Geochem.* 31 (2000) 945–958.

[31] J. Kaal, A. Martínez Cortizas, K.G.J Nierop,. Characterisation of aged charcoal using a coil probe pyrolysis-GC/MS method optimised for black carbon. *J. Anal. Appl. Pyrolysis* 85 (2009) 408–416.

[32] J.A. Baldock, R.J. Smernik, Chemical composition and bioavailability of thermally, altered *Pinus resinosa* (Red Pine) wood. *Org. Geochem.* 34 (2002) 1093–1109.

[33] M.W.I. Schmidt, A.G. Noack, Black carbon in soils and sediments: Analysis, distribution, implications, and current challenges. *Global Biogeochem. Cy.* 14 (2000) 777–793.

[34] N. Poirier, S. Derenne, J. Balesdent, J.-N. Rouzaud, A. Mariotti, C. Largeau, Abundance and composition of the refractory organic fraction of an ancient, tropical soil (Pointe Noire, Congo). *Org. Geochem.* 33 (2002) 383–391.

[35] N. Poirier, S. Derenne, J.-N. Rouzaud, C. Largeau, A. Mariotti, J. Balesdent, J. Maquet, Chemical structure and sources of the macromolecular, resistant, organic

fraction isolated from a forest soil (Lacadee, South-West France). *Org. Geochem.* 31 (2000) 813–827.

[36] K. Quénéa, S. Derenne, F.J. González-Vila, A. Mariotti, J.-N. Rouzaud, C.

Largeau, Study of the composition of the macromolecular refractory fraction from an acidic sandy forest soil (Landes de Gascogne, France) using chemical degradation and electron microscopy. *Org. Geochem.* 36 (2005) 1151–1162.

[37] J. Kaal, A. Martínez Cortizas, E. Meckmeier, M. Costa Casais, M. Santos

Estévez, F. Criado Boado, Holocene fire history of black colluvial soils revealed by pyrolysis-GC/MS: a case study from Campo Lameiro (NW Spain). *J. Archeol. Sci.* 35 (2008) 2133–2143.

[38] G. Almendros, H. Knicker, F.J. González-Vila, Rearrangement of carbon and nitrogen forms in peat after progressive thermal oxidation as determined by solid-state C^{13} and N^{15} -NMR spectroscopy. *Org. Geochem.* 34 (2003) 1559–1568.

[39] Ö. Gustafsson, P.M. Gschwend, The flux of black carbon to surface sediments on the New England continental shelf. *Geochim. Cosmochim. Acta.* 62 (1998) 465–472.

[40] J. Kaal, A. Martínez-Cortizas, K.G.J. Nierop, P. Buurman, A detailed pyrolysis-GC/MS analysis of a black carbon-rich acidic colluvial soils (Atlantic Ranker) from NW Spain. *Appl. Geochem.* 23 (2008) 2395–2405.

[41] G. Cornelissen, J. Haftka, J. Parsons, Ö. Gustafsson, Sorption to black carbon of organic compounds with varying polarity and planarity. *Environ. Sci. Technol.* 39 (2005) 3688–3694.

[42] Y. Zhang, S. Tao, Global atmospheric emission inventory of polycyclic aromatic hydrocarbons (PAHs) for 2004. *Atmos. Environ.* 43 (2009) 812–819

455 [43] S.C. Brassell, G. Eglinton, Environmental chemistry—an interdisciplinary
456 subject. Natural and pollutant organic compounds in contemporary aquatic
457 ecosystems, In: J. Albaigés (Ed) Analytical Techniques in Environmental
458 Chemistry, Pergamon Press, Oxford, 1980.

459 [44] H.H. Soclo, P.H. Garrigues, M. Ewald (2000) Origin of polycyclic aromatic
460 hydrocarbons (PAHs) in coastal marine sediments: Case studies in Cotonou (Benin)
461 and Aquitaine (France) areas. Mar. Pollut. Bull. 40 (2000) 387–396.

462 [45] S. Killops, V. Killops, Introduction to Organic Geochemistry (2nd Ed.) Blackwell
463 Publishing Ltd. 2005.

464 [46] H. Knicker, K.U. Totsche, G. Almendros, F.J. González-Vila, Condensation
465 degree of burnt peat and plant residues and the reliability of solid-state VACP MAS
466 ¹³C NMR spectra obtained from pyrogenic humic material. Org. Geochem. 36
467 (2005) 1359–1377.

468 [47] D. Fabbri, A.G. Rombolá, C. Torri, K.A. Spokas, Determination of polycyclic
469 aromatic hydrocarbons in biochar and biochar amended soil. J. Anal. Appl.
470 Pyrolysis 103 (2013) 60–67.

471 [48] F.J. González-Vila, J.L. López, F. Martin, J.C. del Río, Determination in soils of
472 PAH produced by combustion of biomass under different conditions. Fresen. J.
473 Anal. Chem. 339 (1991) 750–753.

474 [49] C. Jiang, R. Alexander, R.I. Kagi, A.P. Murray, Polycyclic aromatic
475 hydrocarbons in ancient sediments and their relationships to palaeoclimate. Org.
476 Geochem. 29 (1998) 1721–1735

477 [50] M. Medina-Vera, Pyrolysis-gas chromatography/mass spectrometry used for
478 screening polycyclic aromatic hydrocarbons by desorption from sediment. J. Anal.
479 Appl. Pyrolysis 36 (1996) 27–35.

480 [51] P. Faure, P. Landais, Rapid contamination screening of river sediments by flash
 481 pyrolysis-gas chromatography–mass spectrometry (PyGC–MS) and
 482 thermodesorption GC–MS (TdGC–MS). *J. Anal. Appl. Pyrolysis* 57 (2001)
 483 187–202.

484 [52] P. Faure, L. Schlepp, L. Mansuy-Huault, M. Elie, E. Jardé, M. Pelletier,
 485 Aromatization of organic matter induced by the presence of clays during flash
 486 pyrolysis-gas chromatography–mass spectrometry (PyGC–MS) A major analytical
 487 artefact. *J. Anal. Appl. Pyrolysis* 75 (2006) 1–10.

488 [53] M. Ochsenkühn-Petropoulou, K. Staikos, G. Matuschek, A. Kettrup, On-line
 489 determination of polycyclic aromatic hydrocarbons in airborne particulate matter by
 490 using pyrolysis/GC–MS. *J. Anal. Appl. Pyrolysis* 70 (2003) 73–85.

491 [54] M. Nali, F. Corana, L. Montanari, L. Pellegrini, A pyrolysis-gas chromatography
 492 mass-spectrometry study on coals. *J. Anal. Appl. Pyrolysis* 29 (1994) 15–23.

493 [55] H. Knicker, How does fire affect the nature and stability of soil organic nitrogen
 494 and carbon? A review. *Biogeochemistry* 85 (2007) 91–118.

495 [56] J.A. González-Pérez, A. Chabbi, J.M. de la Rosa Arranz, C. Rumpel, F.J.
 496 González-Vila, Evolution of organic matter in lignite-containing sediments revealed
 497 by analytical pyrolysis (Py-GC/MS). *Organic Geochemistry* 53 (2012) 119–130.

498 [57] B.R.T. Simoneit, Cyclic terpenoids of the geosphere, In: R.B. Johns (Ed.),
 499 *Biological Markers in the Sedimentary Record*. Elsevier, Amsterdam, 1986.

500 [58] M. Elmquist, G. Cornelissen, Z. Kukulska, Ö. Gustafsson, Distinct oxidative
 501 stabilities of char versus soot black carbon: Implications for quantification and
 502 environmental recalcitrance, *Global Biogeochem. Cy.* 20 (2006) GB2009 (DOI:
 503 10.1029/2005GB002629).

504 [59] K. Hammes, M.W.I. Schmidt, R.J. Smernik, L.A.R. Currie, W.P. Ball, T.H.
505 Nguyen, P. Louchouart, S. Houel, Ö. Gustafsson, M. Elmquist, G. Cornelissen, J.O.
506 Skjemstad, C.A. Masiello, J. Song, P. Peng, S. Mitra, J.C. Dunn, P.G. Hatcher, W.C.
507 Hockaday, D.M. Smith, C. Hartkopf-Fröder, A. Böhmer, B. Luer, B.J. Huebert, W.
508 Amelung, S. Brodowski, L. Huang, W. Zhang, P.M. Gschwend, D.X. Flores-
509 Cervantes, C. Largeau, J.-N. Rouzaud, C. Rumpel, G. Guggenberger, K. Kaiser, A.
510 Rodionov, F.J. González-Vila, J.A. González-Pérez, J.M. de la Rosa, D.A.C.
511 Manning, E. López-Capél, L. Ding, Comparison of black carbon quantification
512 methods to measure fire-derived (black/elemental) carbon in soils and sediments
513 using reference materials from soil, water, sediment and the atmosphere. *Global*
514 *Biogeochem. Cy.* 21 (2007) GB3016 (DOI: 10.1029/2006GB002914).

515 [60] J.A. Vera, Evolution of the South Iberian Continental Margin. *Memoir. Mus.*
516 *Natl. Hist. Nat.* 186 (2001) 109–143.

517 [61] F. García-García, G.A. De Gea, P.A. Ruiz-Ortiz, Detached forced-regressive
518 shoreface wedges at the Southern Iberian continental palaeomargin (Early
519 Cretaceous, Betic Cordillera, S. Spain). *Sedimentary Geology* 236 (2011) 197–210.

520 [62] J.M. de la Rosa, J.A. González-Pérez, P.G. Hatcher, H. Knicker, F.J. González-
521 Vila, Determination of refractory organic matter in marine sediments by chemical
522 oxidation, analytical pyrolysis and solid-state ¹³C nuclear magnetic resonance
523 spectroscopy. *Eur. J. Soil. Sci.* 59 (2008) 430–438.

524 [63] P. Morf, P. Hasler, T. Nussbaumer, Mechanisms and kinetics of homogeneous
525 secondary reactions of tar from continuous pyrolysis of wood chips. *Fuel* 81 (2002)
526 843–853.

527 [64] C.M. Lakshmanan, H.E. Hoelscher, Production of levoglucosan by pyrolysis of
528 carbohydrates - pyrolysis in hot inert gas stream. Ind. Eng. Chem. Prod. Res. Dev. 9
529 (1970) 57–79.

530 [65] M.B. Yunker, R.W. Macdonald, R. Vingarzan, R.H. Mitchell, D. Goyette, S.
531 Sylvestre, PAHs in the Fraser River basin: a critical appraisal of PAH ratios as
532 indicators of PAH source and composition. Org. Geochem. 33 (2002) 489–515.

533 [66] W.J. Bond, A.C. Scott, Fire and the spread of flowering plants in the Cretaceous.
534 New Phytol. 188 (2010) 1137–1150.

535 [67] J.M. Moreno, W.C. Oechel, The Role of Fire in Mediterranean-type Ecosystems.
536 Ecological Studies Vol. 107. Springer, New York, 1995.

537 [68] L Trabaud, Man and fire: Impacts on Mediterranean vegetation. In: F. di Castri,
538 D.W. Goodall and R.L. Spetch (Eds.). Ecosystems of the World, vol. 11. Elsevier,
539 Amsterdam, 1984.

540 [69] Z. Naveh, Fire in the Mediterranean: A landscape ecological perspective. In:
541 J.G. Goldammer and M.J. Jenkins editors. Fire in ecosystems dynamics:
542 Mediterranean and Northern perspective. SPB Academic Publishing, The Hague,
543 1990.

544 [70] M. Barberó, R. Loisel, P. Quezel, Signification biogéographique et biodiversité
545 des forêts du bassin méditerranéen In: D.M. Richardson (Ed.), Ecology and
546 Biogeography of *Pinus*. Cambridge Univ. Press, Cambridge, 1998.

547 [71] J. Banthus, P.K. Khanna, R.J. Raison, The effect of fire on carbon and nitrogen
548 mineralization and nitrification in an Australian forest soil. Aust. J. Soil. Res. 31
549 (1993) 621–639.

550 [72] J. Pietikäinen, H. Fritze, Clear-cutting and prescribed burning in coniferous
551 forest - comparison of effects on soil fungal and total microbial biomass, respiration
552 activity and nitrification. *Soil. Biol. Biochem.* 27 (1995) 101–109.

553 [73] M.E. Arias, J.A. González-Pérez, F.J. González-Vila, A.S. Ball, Soil health - a
554 new challenge for microbiologists and chemists. *Int. Microbiol.* 8 (2005) 13–21.

555 [74] P.J. Viro, Effect of forest fire on soil. In: C.E. Kozlowski, C.E. Ahlgren, editors.
556 Fire and ecosystems. Academic Press, New York, 1974.

557

TABLE CAPTIONS

Table 1. Materials used

FIGURE CAPTIONS

Fig. 1. Pyrograms at 800 °C of selected reference materials: SIM traces for pyrogenic PAHs: m/z 178 (phenanthrene/anthracene); 202 (fluorene/pyrene); 228 (benzanthracene/chrysene); 252 (benzofluorene/benzopyrene); 276 (indenopyrene/benzoperylene) and m/z 85 for *n*-alkanes.

Fig. 2. Solid-state ^{13}C NMR spectra and Py-GC/MS total current ion pyrograms of BC sample isolated from a marine sediment (Huelva, Spain) and charred chestnut wood.


Fig.3. Series of PAHs identified by analytical pyrolysis (500 °C) of Cretaceous sediment (shale) from the Cerrajón Formation (SE Iberian Peninsula).


Fig. 4. Analytical pyrolysis (500 °C) of wildfire-affected and control, not affected soils from “Sierra de Cazorla” (S. Iberian Peninsula). Ion chromatogram traces (m/z 91 alkylbenzenes; m/z 178 three ring PAHs, m/z 228 four ring PAHs and m/z 252 five ring PAHs).

Table 1. Materials analysed





Material	Production/collection method	Rationale	Selected reference
Group A: Natural samples and laboratory produced PAH-rich materials			
Soot: from <i>n</i> -hexane	Collected from the top of an <i>n</i> -hexane flame	Condensate produced from a fossil fuel	[52]
Char: Chestnut wood	<i>Castanea sativa</i> wood at 400 °C for 5 h in N atmosphere	Char produced from hardwood biomass	[51]
Char: Rice grass	<i>Oryza sativa</i> straw at 400 °C for 5 h in N atmosphere	Char produced from grass biomass	[51]
Coal: Low bituminous	From Buchanan County (Virginia, USA)	A mature coal (c. 10% pyrogenic inertinite)	[52]
Group B: Environmental matrices			
Soil: Vertisol	Sandy clay soil from Urrbrae (Australia)	PAHs in soils	[51]
Soil: Chernozem	Light sandy soil from Harsum (Germany)	PAHs in soils	[51]
Soil: Rendollic soil	Soil on loess from Sierra de Cazorla (Jaén, Spain)	Pyrogenic PAHs from wildfire-affected soil (2005)	[12]
Sediment: Turbidite	Early Cretaceous formation from the Cerrajón (Jaén, Spain)	Turbidite with fossil pyrogenic PAHs	[56–57]
Group C: Negative controls that contains no or low PAH content			
Melanoidin	Produced by heating urea and glucose at 90 °C for 30 days	Negative control, contains no PAHs	[51]
ROM* sediment fraction	From offshore marine sediment (Southwest Spain)	Negative control, contains no PAHs	[58]





*) Organic matter refractory to wet oxidation at room temperature





○ *n*-alkanes ● *n*-fatty acids □ Benzofuran derivatives ◆ Aromatic structures ◆ PAHs ★ Anhydrosugars





Traces are normalised with that of *n*-heptylbenzene (+)

